

Introduction to lifecontingencies Package

Giorgio Alfredo Spedicato, Ph.D

Abstract

lifecontingencies performs actuarial present value calculation for life insurances. This paper briefly recapitulate the theory regarding life contingencies (life tables, financial mathematics and related probabilities) on life contingencies. Then it shows how **lifecontingencies** functions represent a perfect cookbook to perform life insurance actuarial analysis and related stochastic simulations.

Keywords: life tables, financial mathematics, actuarial mathematics, life insurance, R.

1. Introduction

As of February 2012, **lifecontingencies** appears to be the first R package that deals with life insurance evaluation.

Some actuarial packages have been already available in R, however most of these packages mainly interest non-life actuaries.

In fact non - life insurance modeling uses more data analysis and applied statistical modelling than life insurance does. E.g. functions to fit loss distributions and to perform credibility analysis are provide within the package **actuar**, Dutang, Goulet, and Pigeon (2008). Package **actuar** represents the computational side of the classical actuarial manual Loss Distribution, Klugman, Panjer, Willmot, and Venter (2009). The package **ChainLadder**, Gesmann and Zhang (2011), provides functions to estimate non-life unpaid loss reserve. GLM models, widely used in non - life insurance pricing, can be fit by functions bundled in the base R distribution. More advanced predictive models used by actuaries, e.g. GAMLSS and Tweedie regressions, can be fit using specifically developed packages as **gamlss**, Rigby and Stasinopoulos (2005), and **cplm** packages respectively.

Life insurance evaluation models demographic and financial data mainly . R has a dedicated view to packages specifically tailored to financial analysis. But, few packages that handle demographic data have been published yet. Relevant packages that can aid demographers work are **demography**, Rob J Hyndman, Heather Booth, Leonie Tickle, and John Maindonald (2011), and **LifeTables**, Riffe (2011). Packages **YieldCurve**, Guirreri (2010), and **termstrc**, Ferstl and Hayden (2010), can be used to perform interest rate analysis. Finally no package exists that performs life contingencies calculations, as of February 2012.

Numerous commercial software specifically tailored to actuarial analysis are available in commerce. Moses and Prophet are currently the leading actuarial software for life insurance modelling. **lifecontingencies** package aims to represent the R computational side of the concepts exposed in the classical SOA Actuarial Mathematics book, Bowers, Gerber, Hickman,

Jones, and Nesbitt (1997). Since life contingencies theory grounds on demography and classical financial mathematics, I have made use of the Ruckman and Francis Ruckman and Francis and Broverman Broverman (2008) as references. The structure of the vignette document is:

1. Section 2 outlines the statistical and financial mathematics theory regarding life contingencies.
2. Section 3 overviews the structure of the **lifecontingencies** package.
3. Section 4 gives a wide choice of applied **lifecontingencies** examples .
4. Finally Section 5 discusses package actual and prospective development and known limitations.

2. Life contingencies statistical and financial foundations

Life insurance analysis involves the calculation of statistics regarding occurrence and amount of future cash flows. E.g. the insurance pure premium (also known as benefit premium) is the present value of the series of future cash flows whose probability is based on the occurrence of the policyholder's life events (life contingencies). Therefore life insurance actuarial mathematics grounds itself on concepts derived from demography and the theory of interest.

A life table (also called a mortality table or actuarial table) is a table that shows how mortality affects subject of a cohort across different ages. It reports for each age x , the number of subjects l_x living at the beginning of age x . It represents a sequence of $l_0, l_1, \dots, l_\omega$, where ω is the farthest age until which a subject of the cohort can survive. Life table are typically distinguished according to gender, year of birth and nationality.

Using a statistical perspective, a life table allows the probability distribution of the the future lifetime for a subject aged x , to be deduced. In particular a life table allows to derive two key probability distribution: T_x , the future lifetime for a subject aged x and its curtate form, K_x , i.e., the number of future years completed before death. Therefore many statistics can be derived from the life table. A non exhaustive list follows:

- ${}_tp_x = \frac{l_{x+t}}{l_x}$, the probability that someone living at age x will reach age $x + t$.
- ${}_tq_x$, the complementary probability of ${}_tp_x$.
- ${}_td_x$, the number of deaths between age x and $x + t$.
- ${}_tL_x = \sum_{t=0}^n l_{x+t}$, the expected number of years lived by the cohort between ages x and $x + t$.
- ${}_tm_x = \frac{{}_td_x}{{}_tL_x}$, the central mortality rate between ages x and $x + t$.
- e_x , the curtate expectation of life for a subject aged x , $e_x = E(K_x)$ and its complete form $\overset{\circ}{e}_x = E(T_x)$.

The Keyfitz manual, [Keyfitz and Caswell \(2005\)](#), provides an exhaustive coverage about life table theory and practice. Life table are usually published by institutions that have access to large amount of reliable historical data, like government statistics or social security bureaus. It is a common practice for actuaries to start from these life tables and to adapt them to the insurer's portfolio actual experience.

Classical financial mathematics deals with monetary amount that could be available in different times. The present value of a series of cash flows, reported in formula 2, is probably the most important concept. The present value represents the current value of a series of monetary cash flows, CF_t , that will be available in different periods of time.

The interest rates, i_t , represents the measure of the price of money available in future times. This paper will use i to express the effective (real) compound interest. It means that if i is the interest rate, a sum of 1 monetary unit accumulates through time according to the law $A(t) = (1 + i)^t$, being $A(t)$ the accumulation function. Arrangements lead to discount and nominal (m-compound) interest rates as shown by equation 1.

$$A(t) = (1 + i)^t = (1 - d)^{-t} = \left(1 + \frac{i^m}{m}\right)^{t*m} = \left(1 - \frac{d^m}{m}\right)^{-t*m} \quad (1)$$

All financial mathematics functions (as annuities, $\bar{a}_{\overline{n}|}$, or accumulated values, $s_{\overline{n}|}$) can be written as a particular case of formula 2. See the classical [Broverman \(2008\)](#) manual for further reference on the topic.

$$PV = \sum_{t \in T} CF_t (1 + i_t)^{-t} \quad (2)$$

Actuaries uses the probabilities inherent the life table to evaluate life contingencies insurances. Life contingencies are themselves stochastic variables, in fact. They consist in present values whose amounts are not certain, since the time of their occurrence and the final values depend by events regarding the life of the insured head. **lifecontingencies** package provides function to model many of such random variables, \tilde{Z} , and in particular their expected value, the Actuarial Present Value (APV). APV is certainly the most important statistic of \tilde{Z} variables that actuaries use. It represents the average cost of the benefit the insurer provides to the policyholder. Insured benefits and loadings for expense and profit adds to the final premium proposed to policyholders. Life contingencies can be either continue or discrete. **lifecontingencies** package models only discrete life contingencies, that is insured amounts are supposed to be due at the end of each year or fraction of year. The ? manual contains formulas to obtain continue life contingencies APV from the corresponding discrete forms.

Few examples of life contingencies follow:

1. An n-year term life insurance provides payment of b if the insured dies within n years from issue. If the payment is performed at the end of year of death, we can write \tilde{Z} as
$$\tilde{Z} = \begin{cases} b * v^{\tilde{K}_x+1}, & \tilde{K}_x \leq n \\ 0, & \tilde{K}_x > n \end{cases}$$
 The APV symbol is $A_{x:\overline{n}|}^1$.
2. A life annuity consists in a series of benefits paid contingent upon survival of a given life. In particular, a temporary life annuity due pays a benefit at the beginning of each

period so long as the annuitant (x) survives, for up to a total of n years, or n payments. Assuming \$1 payments, we can write \tilde{Z} as $\tilde{Z} = a_{\overline{\tilde{K}+1}|}$. Its APV expression is $\ddot{a}_{x:\overline{n}|}$.

3. An n -year pure endowment insurance grants a benefit payable at the end of n years if the insured survives at least n years from issue. The expression of \tilde{Z} is $v^n * I(\tilde{K}_x \geq n)$ and its APV expression is ${}_nE_x$.

We remained to the [Bowers *et al.* \(1997\)](#) manual for formulas regarding other life contingencies insurances (as $(DA)_{x:\overline{n}|}^1$, the decreasing term life insurance, $(IA)_x$, the increasing term life insurance for example) and common variations on benefit payment: deferral and fractional ages handling.

The **lifecontingencies** package provides functions that allows the actuary to evaluate the APV and to draw random samples from \tilde{Z} distribution. Three approach have been traditionally followed for the evaluation of the APV: the use of commutation tables, the current payment technique and the expected value techniques. Commutation tables extend life table by tabulating special function of age and rate of interest whose ratios allow the actuary to evaluate APV for standard insurances, as discussed in [Anderson, of Actuaries. Education, and Committee \(1999\)](#). The **lifecontingencies** allows underlying commutation table to be printed out as further described. However commutation table usage has become useless in computer era since it does not allows varying benefits payments to be modelled easily and it is computationally inefficient. Therefore commutation table approach has not been used within **lifecontingencies** to perform APV calculations.

The current payment technique returns the APV calculation as the scalar product of three vectors, as formula 3 shows: the vector of uncertain cash flows, \bar{c} , the vector of discount factors, \bar{v} and the vector of cash flow probability. Since this approach is the most computationally efficient we have used this approach to evaluate APV. The expected value approach models the APV as the scalar product of two vector: $\Pr[\tilde{K} = k]$, the probability that the future curtate lifetime is exactly k , where $\tilde{K}_x = 0, \dots, \omega - x$, the present value of the benefit due under the insurance terms if the future curtate lifetime is exactly k . The latter approach has been used to define the probability distribution of \tilde{Z} in order to allow random sampling from its distribution.

$$\langle \langle \bar{c} \bullet \bar{v} \rangle \bullet \bar{p} \rangle \quad (3)$$

3. The structure of the package

Package **lifecontingencies** contains classes and methods to handle lifetables and actuarial tables conveniently.

The package is loaded within the R command line as follows:

```
R> library(lifecontingencies)
```

Two main S4 classes [Chambers \(2008\)](#) have been defined within the **lifecontingencies** package: the `lifetable` class and the `actuarialtable` class. The `lifetable` class is defined as follows

```
R> #definition of lifetable
R> showClass("lifetable")
```

```
Class "lifetable" [in ".GlobalEnv"]
```

```
Slots:
```

```
Name:      x      lx      name
Class:  numeric numeric character
```

```
Known Subclasses: "actuarialtable"
```

Class `actuarialtable` inherits from `lifetable` class and has another additional slots, the interest rate.

```
R> showClass("actuarialtable")
```

```
Class "actuarialtable" [in ".GlobalEnv"]
```

```
Slots:
```

```
Name:  interest      x      lx      name
Class:  numeric  numeric numeric character
```

```
Extends: "lifetable"
```

Beyond generic S4 classes and methods there are three groups of functions: demographics, financial mathematics and life contingencies analysis functions.

The demographic group comprises the following functions:

1. `dxt` returns deaths between age x and $x + t$, $d_{x,t}$.
2. `pxt` returns survival probability between age x and $x + t$, $p_{x,t}$.

3. **pxyt** returns the survival probability for two lives, $d_{xy,t}$.
4. **qxt** returns death probability between age x and $x + t$, $q_{x,t}$.
5. **qxyt** returns the survival probability for two lives, $q_{xy,t}$.
6. **Txt** returns the number of person-years lived after exact age x , $T_{x,t}$.
7. **mxt** returns central mortality rate, $m_{x,t}$.
8. **exn** returns the complete or curtate expectation of life from age x to $x + n$, $e_{x,n}$.
9. **rLife** returns a sample from the time until death distribution underlying a life table.
10. **exyt** returns the expected life time for two lives between age x and $x + t$.
11. **probs2lifetable** returns a life table l_x from raw one - year survival / death probabilities.

The financial mathematics group comprises the following functions, for which we report most important function:

1. **presentValue** returns the present value for a series of cash flows.
2. **annuity** returns the present value of a annuity - certain, $a_{\overline{n}|}$.
3. **increasingAnnuity** returns the present value of an increasing annuity - certain, $(IA)_n$.
4. **accumulatedValue** returns the future value of a series of cash flows, $s_{\overline{n}|}$.
5. **decreasingAnnuity** returns the present value of an increasing annuity, $(DA)_{\overline{n}|}$.
6. **accumulatedValue** returns the future value of a payments sequence, $s_{\overline{n}|}$.
7. **nominal2Real** returns the effective annual interest (discount) rate i given the nominal m-periodal interest $i^{(m)}$ or discount d^m rate.
8. **real2Nominal** returns the m-periodal interest or discount rate given the m periods or the discount.
9. **intensity2Interest** returns the intensity of interest δ given the interest rate i .
10. **interest2Intensity** returns the interest rate i given the intensity of interest δ .

The actuarial mathematics group comprises the following functions, for which we report must important function:

1. **Axn** returns the APV for life insurances.
2. **Axyn** returns the APV for two heads life insurances.
3. **axn** returns the APV for annuities.
4. **axyn** returns the APV for two heads annuities.

5. **Exn** returns the APV for the pure endowment.
6. **Iaxn** returns the APV for the increasing annuity.
7. **IAxn** returns the APV for the increasing life insurance.
8. **DAxn** returns the APV for the decreasing life insurance.

Draft

4. Code and examples

4.1. Classical financial mathematics example

The **lifecontingencies** package provides functions to perform classical financial analysis. Following examples will show how to handle interest and discount rates with different compounding frequency, how to perform present value, annuities and future values analysis calculations, loans amortization and bond pricing.

Interest rate functions

Following examples show how to switch from $i^m \rightarrow i$

```
R> #an APR of 3% is equal to a
R> real2Nominal(0.03,12)
```

```
[1] 0.02959524
```

```
R> #of nominal interest rate while
R> #6% annual nominal interest rate is the same of
R> nominal2Real(0.06,12)
```

```
[1] 0.06167781
```

```
R> #APR
R> #4% per year compounded quarterly is
R> nominal2Real(0.04,4)
```

```
[1] 0.04060401
```

```
R> #4% effective interest rate corresponds to
R> real2Nominal(0.04,4)*100
```

```
[1] 3.941363
```

```
R> #nominal interest rate (in 100s) compounded quarterly
```

and from $d^m \rightarrow d$

```
R> #a nominal rate of discount of 4% payable quarterly is equal to a
R> real2Nominal(i=0.04,m=12,type="discount")
```

```
[1] 0.04075264
```

Present value analysis

Performing a project appraisal means evaluating the present value of all net cash flows, as shown in code below:


```
R> #suppose an investment requires and grants following cash flows
R> capitals=c(-1000,200,500,700)
R> #at time (vector) t.
R> times=c(-2,-1,4,7)
R> #the preset value of the investment is
R> presentValue(cashFlows=capitals, timeIds=times,
+               interestRates=0.03)
```

```
[1] 158.5076
```

```
R> #assuming 3% interest rate
R>
R> #while if interest rates were time - varying
R> #e.g. 0.04 0.02 0.03 0.057
R> presentValue(cashFlows=capitals, timeIds=times,
+               interestRates=c( 0.04, 0.02, 0.03, 0.057))
```

```
[1] 41.51177
```

```
R> #and if the last cash flow is uncertain, as we assume a
R> #receiving probability of 50%
R> presentValue(cashFlows=capitals, timeIds=times,
+               interestRates=c( 0.04, 0.02, 0.03, 0.057),
+               probabilities=c(1,1,1,0.5))
```

```
[1] -195.9224
```

Annuities and future values

Example of $a_{\overline{n}|}$ and $s_{\overline{n}|}$ evaluations are reported below.

```
R> #PV annuity immediate 100$ each year 5 years @9%
R> 100*annuity(i=0.09,n=5)
```

```
[1] 388.9651
```

```
R> #while the corresponding future values is
R> 100*accumulatedValue(i=0.09,n=5)
```

```
[1] 598.4711
```

```
R> #A man wants to save 100,000 to pay for the education
R> #of his son in 10 years time. An education fund requires the investors to
R> #deposit equal instalments annually at the end of each year. If interest of
R> #0.075 is paid, how much does the man need to save each year (R) in order to
R> #meet his target?
R> 100000/accumulatedValue(i=0.075,n=10)
```

```
[1] 7068.593
```

while the code below shows how fractional annuities ($a_{\overline{t}|i}^{(m)}$) can be handled within `annuity` and `accumulatedValue` functions.

```
R> #Find the present value of an annuity-immediate of
R> #100 per quarter for 4 years, if interest is compounded semiannually at
R> #the nominal rate of 6%.
R> #the APR is
R> APR=nominal2Real(0.06,2)
R> 100*4*annuity(i=APR,n=4,m=4)
```

```
[1] 1414.39
```

Finally `increasingAnnuity` and `decreasingAnnuity` functions handle increasing $((IA)_x)$ and decreasing $((DA)_x)$ annuities.

```
R> #An increasing n-payment annuity-due shows payments of 1, 2,
R> #... , n
R> #at time 0, 1, ... ,
R> #n - 1 . At interest rate of
R> #0.03 and n=10, its present value of the annuity is
R> increasingAnnuity(i=0.03, n=10,type="due")
```

```
[1] 46.18416
```

```
R> #while the present value of a decreasing
R> #annuity due of 10, 9,...,1
R> #from time 1 to time 10 is
R> decreasingAnnuity(i=0.03, n=10,type="immediate")
```

```
[1] 48.99324
```

Finally the calculation of the present value of a geometrically increasing annuity is shown in the code below

```
R> #assume each year the annuity increases its value by 3%
R> #while the interest rate is 4%
R> #first determine the effective interest rate
R> ieff=(1+0.04)/(1+0.03)-1
R> #assume the annuity lasts 10 years
R> annuity(i=ieff,n=10)
```

```
[1] 9.48612
```

Loan amortization

The code lines below show how an investment amortization schedule will be repaired.

Suppose loaned capital is C , then assuming an interest rate i , the amount due to the lender at each instalment is $R = \frac{C}{a_{\overline{n}|i}}$.

At each installment the R_t installment repays $I_t = C_{t-1} * i$ as interest and $C_t = R_t - I_t$ as capital.

```
R> capital=100000
R> interest=0.05 #assume 5% effective annual interest
R> payments_per_year=2 #payments per year
R> rate_per_period=(1+interest)^(1/payments_per_year)-1
R> years=5 #five years length of the loan
R> installment=1/payments_per_year*capital/annuity(i=interest, n=years,m=payments_per_year)
R> installment
```

```
[1] 11407.88
```

```
R> #compute the balance due at the begin of period
R> balance_due=numeric(years*payments_per_year)
R> balance_due[1]=capital*(1+rate_per_period)-installment
R> for(i in 2:length(balance_due))
+ {
+     balance_due[i]=balance_due[i-1]*(1+rate_per_period)-installment
+     cat("Payment ",i, " balance due:",round(balance_due[i]),"\n")
+ }
```

```
Payment 2 balance due: 81903
Payment 3 balance due: 72517
Payment 4 balance due: 62900
Payment 5 balance due: 53046
Payment 6 balance due: 42948
Payment 7 balance due: 32600
Payment 8 balance due: 21998
Payment 9 balance due: 11133
Payment 10 balance due: 0
```

Bond pricing

Bond pricing is another application of present value analysis. A standard bond whose principal will be repaid at time T is a series of coupon c_t , priced according to a coupon rate $j^{(k)}$ on a principal C . Formula 4 expresses the present value of a bond.

$$B_t = c_t a_{\overline{n}|i}^{(k)} + C v^T \quad (4)$$

We will show how to evaluate a standard bond with following examples:

```
R> bond<-function(faceValue, couponRate, couponsPerYear, yield,maturity)
+ {
+     out=NULL
+     numberOfCF=maturity*couponsPerYear #determine the number of CF
+     CFs=numeric(numberOfCF)
+     payments=couponRate*faceValue/couponsPerYear #determine the coupon sum
+     cf=payments*rep(1,numberOfCF)
+     cf[numberOfCF]=faceValue+payments #set the last payment amount
+     times=seq.int(from=1/couponsPerYear, to=maturity, by=maturity/numberOfCF)
+     out=presentValue(cashFlows=cf, interestRates=yield, timeIds=times)
+     return(out)
+ }
R> #bond coupon rate 6%, two coupons per year, face value 1000, yield 5%, three years to m
R> bond(1000,0.06,2,0.05,3)

[1] 1029.25

R> #bond coupon rate 3%, one coupons per year, face value 1000, yield 3%, three years to m
R> bond(1000,0.06,1,0.06,3)

[1] 1000
```

4.2. Lifetables and actuarial tables analysis

`lifetable` classes represent the basic class designed to handle life table calculations. A `actuarialtable` class inherits from `lifetable` class adding one more slot to set the a priori rate of interest.

Both classes have been designed using the S4 class framework.

Examples follow showing how `lifetable` and `actuarialtable` objects initialization, basic survival probability and life tables analysis.

Creating lifetable and actuarialtable objects

Lifetable objects can be created by raw R commands or using existing `data.frame` objects. However, to build a `lifetable` class object three items are needed:

1. The years sequence, that is an integer sequence $0, 1, \dots, \omega$. It shall starts from zero and going to the ω age (the age x that $p_x = 0$).
2. The l_x vector, that is the number of subjects living at the beginning of age x .
3. The name of the life table.

```
R> x_example=seq(from=0,to=9, by=1)
R> lx_example=c(1000,950,850,700,680,600,550,400,200,50)
R> fakeLt=new("lifetable",x=x_example, lx=lx_example, name="fake lifetable")
```

A `print` (or `show`) method are available. These methods report the x , l_x , p_x and e_x in tabular form.

```
R> print(fakeLt)
```

Life table fake lifetable

	x	lx	px	ex
1	0	1000	0.9500000	4.742105
2	1	950	0.8947368	4.241176
3	2	850	0.8235294	4.042857
4	3	700	0.9714286	3.147059
5	4	680	0.8823529	2.500000
6	5	600	0.9166667	1.681818
7	6	550	0.7272727	1.125000
8	7	400	0.5000000	0.750000
9	8	200	0.2500000	0.500000

`head` and `tail` methods for `data.frame` S3 classes have also been adapted to `lifetable` classes, as code below shows.

```
R> #show head method
R> head(fakeLt)
```

```

  x   lx
1 0 1000
2 1  950
3 2  850
4 3  700
5 4  680
6 5  600

```

```

R> #show tail method
R> tail(fakeLt)

```

```

  x   lx
5 4 680
6 5 600
7 6 550
8 7 400
9 8 200
10 9  50

```

Nevertheless the easiest way to create a `lifetable` object is starting from a suitable existing `data.frame`.

```

R> #load USA Social Security LT
R> data(demoUsa)
R> usaMale07=demoUsa[,c("age", "USSF2007M")]
R> usaMale00=demoUsa[,c("age", "USSF2000M")]
R> #coerce from data.frame to lifecontingencies requires x and lx names
R> names(usaMale07)=c("x", "lx")
R> names(usaMale00)=c("x", "lx")
R> #apply coerce methods and changes names
R> usaMale07Lt<-as(usaMale07,"lifetable")
R> usaMale07Lt@name="USA MALES 2007"
R> usaMale00Lt<-as(usaMale00,"lifetable")
R> usaMale00Lt@name="USA MALES 2000"
R> #create the tables
R> ##males
R> lxIPS55M<-with(demoIta, IPS55M)
R> pos2Remove<-which(lxIPS55M %in% c(0,NA))
R> lxIPS55M<-lxIPS55M[-pos2Remove]
R> xIPS55M<-seq(0,length(lxIPS55M)-1,1)
R> ##females
R> lxIPS55F<-with(demoIta, IPS55F)
R> pos2Remove<-which(lxIPS55F %in% c(0,NA))
R> lxIPS55F<-lxIPS55F[-pos2Remove]
R> xIPS55F<-seq(0,length(lxIPS55F)-1,1)
R> #finalize the tables
R> ips55M=new("lifetable",x=xIPS55M, lx=lxIPS55M, name="IPS 55 Males")
R> ips55F=new("lifetable",x=xIPS55F, lx=lxIPS55F, name="IPS 55 Females")

```

The last way a `lifetable` object can be created is generating it from one year survival or death probabilities. Such probabilities could be obtained from mortality projection methods (e.g. Lee - Carter).

```
R>      #use 2002 Italian males life tables
R>      data(demoIta)
R>      itaM2002<-demoIta[,c("X","SIM92")]
R>      names(itaM2002)=c("x","lx")
R>      itaM2002Lt<-as(itaM2002,"lifetable")
```

removing NA and 0s

```
R>      itaM2002Lt@name="IT 2002 Males"
R>      #reconvert in data frame
R>      itaM2002<-as(itaM2002Lt,"data.frame")
R>      #add qx
R>      itaM2002$qx<-1-itaM2002$px
R>      #reduce to 20% one year death probability for ages between 20 and 60
R>      for(i in 20:60) itaM2002$qx[itaM2002$x==i]=0.2*itaM2002$qx[itaM2002$x==i]
R>      #obtain the reduced mortality table
R>      itaM2002reduced<-probs2lifetable(probs=itaM2002[, "qx"], radix=100000,type="qx",
```

An `actuarialtable` class inherits from the `lifecontingencies` class, but it contains an additional slot: the interest rate slot.

```
R>      #assume 3% interest rate
R>      fakeAct=new("actuarialtable",x=fakeLt@x, lx=fakeLt@lx, interest=0.03,
+                  name="fake actuarialtable")
```

Method `getOmega` provides the ω age.

```
R>      getOmega(fakeAct)
```

```
[1] 9
```

Method `print` behaves differently between `lifetable` objects and `actuarialtable` objects. One year survival probability and complete expected remaining life until deaths is reported when `print` method is applied on a `lifetable` object. Classical commutation functions (D_x , N_x , C_x , M_x , R_x) are reported when `print` method is applied on an `actuarialtable` object.

```
R> #apply method print applied on a life table
R> print(fakeLt)
```

Life table fake lifetable

	x	lx	px	ex
1	0	1000	0.9500000	4.742105
2	1	950	0.8947368	4.241176
3	2	850	0.8235294	4.042857
4	3	700	0.9714286	3.147059
5	4	680	0.8823529	2.500000
6	5	600	0.9166667	1.681818
7	6	550	0.7272727	1.125000
8	7	400	0.5000000	0.750000
9	8	200	0.2500000	0.500000

```
R> #apply method print applied on an actuarial table
R> print(fakeAct)
```

Actuarial table fake actuarialtable interest rate 3 %

	x	lx	Dx	Nx	Cx	Mx	Rx
1	0	1000	1000.00000	5467.92787	48.54369	840.7400	4839.7548
2	1	950	922.33010	4467.92787	94.25959	792.1963	3999.0148
3	2	850	801.20652	3545.59778	137.27125	697.9367	3206.8185
4	3	700	640.59916	2744.39125	17.76974	560.6654	2508.8819
5	4	680	604.17119	2103.79209	69.00870	542.8957	1948.2164
6	5	600	517.56527	1499.62090	41.87421	473.8870	1405.3207
7	6	550	460.61634	982.05563	121.96373	432.0128	931.4337
8	7	400	325.23660	521.43929	157.88185	310.0491	499.4210
9	8	200	157.88185	196.20268	114.96251	152.1672	189.3719
10	9	50	38.32084	38.32084	37.20470	37.2047	37.2047

Basic demographic calculations

Basic probability calculations may be performed on valid lifetable or actuarialtable objects. Below calculations for ${}_t p_x$, ${}_t q_x$ and $\dot{e}_{x:\overline{n}|}$.

```
R> #using ips55M life table
R> #probability to survive one year, being at age 20
R> pxt(ips55M,20,1)
```

```
[1] 0.9995951
```

```
R> #probability to die within two years, being at age 30
R> qxt(ips55M,30,2)
```

```
[1] 0.001332031
```



```
R> #expected life time between 50 and 70 years
R> exn(ips55M, 50,20)
```

```
[1] 19.43322
```

Fractional survival probabilities can also be calculated according with linear interpolation, constant force of mortality and hyperbolic assumption.

```
R> data(soa08Act) #load Society of Actuaries illustrative life table
R> pxt(soa08Act,80,0.5,"linear") #linear interpolation (default)
```

```
[1] 0.9598496
```

```
R> pxt(soa08Act,80,0.5,"constant force") #constant force
```

```
[1] 0.9590094
```

```
R> pxt(soa08Act,80,0.5,"hyperbolic") #hyperbolic Balducci's assumption.
```

```
[1] 0.9581701
```

Analysis of two heads survival probabilities can be performed also, as shown by code below:

```
R> pxxt(fakeLt,fakeLt,x=6, y=7, t=2) #joint survival probability
```

```
[1] 0.04545455
```

```
R> pxxt(fakeLt,fakeLt,x=6, y=7, t=2,status="last") #last survival probability
```

```
[1] 0.4431818
```

```
R> #evaluate the expected joint life time for a couple aged 65 and 63 using Italina IPS55
R> exyt(ips55M, ips55F, x=65,y=63, status="joint")
```

```
[1] 19.1983
```

4.3. Classical actuarial mathematics examples

Classical actuarial mathematics on life contingencies will follow now. We will use the SOA illustrative life table on all following examples.

Life insurance examples

Following examples show the APV (i.e. the lump sum benefit premium) for:

1. 10-year term life insurance for a subject aged 30 assuming 4% interest rate, $A_{30:\overline{10}|}^1$.
2. 10-year term life insurance for a subject aged 30 with benefit payable at the end of month of death at 4% interest rate.
3. whole life insurance for a subject aged 40 assuming 4% interest rate, A_{40} .
4. 5 years deferred 10-years term life insurance for a subject aged 40 assuming 5% interest rate, ${}_{5|10}\bar{A}_{40}$.
5. 5 years annually decreasing term life insurance for a subject aged 50 assuming 6% interest rate, $(DA)_{50:\overline{5}|}^1$.
6. 20 years increasing term life insurance, age 40, $(IA)_{50:\overline{5}|}^1$.

```
R> #The APV of a life insurance for a 10-year term life insurance for an
R> #insured aged 40 @ 4% interest rate is
R> Axn(soa08Act, 30,10,i=0.04)
```

```
[1] 0.01577283
```

```
R> #same as above but payable at the end of month of death
R> Axn(soa08Act, x=30,n=10,i=0.04,k=12)
```

```
[1] 0.01605995
```

```
R> #a whole life for a 40 years old insured at @4% is
R> Axn(soa08Act, 40) #soa08Act has 6% implicit interest rate
```

```
[1] 0.1613242
```

```
R> #a 5-year deferred life insurance, 10 years length, 40 years age, @5% interest rate
R> Axn(soa08Act, x=40,n=10,m=5,i=0.05)
```

```
[1] 0.03298309
```

```
R> #Five years annually decreasing term life insurance, age 50.
R> DAxn(soa08Act, 50,5)
```

```
[1] 0.08575918
```

```
R> #Increasing 20 years term life insurance, age 40
R> IAxn(soa08Act, 40,10)
```

```
[1] 0.1551456
```

while following code evaluates pure endowments APV, ${}_nE_x$, assuming SOA life table at 6% interest rate.

```
R> #evaluate the APV for a n year pure endowment, age x=30, n=35, i=6%
R> Exn(soa08Act, x=30, n=35, i=0.06)
```

```
[1] 0.1031648
```

```
R> #try i=3%
R> Exn(soa08Act, x=30, n=35, i=0.03)
```

```
[1] 0.2817954
```

Life annuities examples

Following examples show annuities APV calculations for

1. annuity immediate for a subject aged 65, a_{65} .
2. annuity due for a subject aged 65, \ddot{a}_{65} .
3. 20 years annuity due with monthly fractional payments of \$1000, $\ddot{a}_{65:\overline{20}|}^{(12)}$.

All examples assume SOA life table at 6% interest rate.

```
R> #assuming insured's age x=65 and SOA illustrative life table @6% hold for all examples
R> #annuity immediate
R> axn(soa08Act, x=65, m=1)
```

```
[1] 8.896928
```

```
R> #annuity due
R> axn(soa08Act, x=65)
```

```
[1] 9.896928
```

```
R> #due with monthly payments of $1000 provision
R> 12*1000*axn(soa08Act, x=65,k=12)
```

```
[1] 113179.1
```

```
R> #due with montly payments of $1000 provision, 20 - years term
R> 12*1000*axn(soa08Act, x=65,k=12, n=20)
```

```
[1] 108223.5
```

```
R> #immediate with monthly payments of 1000 provision, 20 - years term
R> 12*1000*axn(soa08Act, x=65,k=12,n=20,m=1/12)
```

```
[1] 107321.1
```

Benefit premiums examples

lifecontingencies package functions can be used to evaluate benefit premium for life contingencies, using the formula ${}_hP_{x:\overline{n}|}^1 = APV\ddot{a}_{x:\overline{n}|}$.

```
R> data(soa08Act) #use SOA MLC exam illustrative life table
R> #Assume X, aged 30, wishes to buy a 250K 35-years life insurance
R> #premium paid annually for 15 years @2.5%.
R> Pa=100000*Axn(soa08Act, x=30,n=35,i=0.025)/axn(soa08Act, x=30,n=15,i=0.025)
R> Pa

[1] 921.5262

R> #if premium is paid montly
R> Pm=100000*Axn(soa08Act, x=30,n=35,i=0.025)/axn(soa08Act, x=30,n=15,i=0.025,k=12)
R> Pm

[1] 932.9836

R> #level semiannual premium for an endowment insurance of 10000
R> #insured age 50, insurance term is 20 years
R> APV=10000*(Axn(soa08Act,50,20)+Exn(soa08Act,50,20))
R> P=APV/axn(soa08Act,50,20,k=2)
```

Benefit reserves examples

Now we will evaluate the benefit reserve for a 20 year life insurance of 100,000, which benefits payable at the end of year of death, which level benefit premium payable at the beginning of each year. Assume 3% of interest rate and SOA life table to apply.

The benefit premium is P , determined from equation

$$P\ddot{a}_{40:\overline{20}|} = 100000A_{40:\overline{20}|}^1$$

. The benefit reserve is ${}_kV_{40+t:\overline{n-t}|}^1 = 100000A_{40+t:\overline{20-t}|}^1 - P\ddot{a}_{40+t:\overline{20-t}|}$ for $t = 0 \dots 19$.

```
R> P=100000*Axn(soa08Act,x=40,n=20,i=0.03)/axn(soa08Act,x=40,n=20,i=0.03)
R> for(t in 0:19) cat("At time ",t," benefit reserve is ", 100000*Axn(soa08Act,x=40,
```

```
At time 0 benefit reserve is 0
At time 1 benefit reserve is 306.9663
At time 2 benefit reserve is 604.0289
At time 3 benefit reserve is 889.0652
At time 4 benefit reserve is 1159.693
At time 5 benefit reserve is 1413.253
At time 6 benefit reserve is 1646.808
At time 7 benefit reserve is 1857.044
```

At time 8 benefit reserve is 2040.286
 At time 9 benefit reserve is 2192.436
 At time 10 benefit reserve is 2308.88
 At time 11 benefit reserve is 2384.513
 At time 12 benefit reserve is 2413.576
 At time 13 benefit reserve is 2389.633
 At time 14 benefit reserve is 2305.464
 At time 15 benefit reserve is 2152.963
 At time 16 benefit reserve is 1922.973
 At time 17 benefit reserve is 1605.162
 At time 18 benefit reserve is 1187.872
 At time 19 benefit reserve is 657.8482

The benefit reserve for a whole life annuity with level annual premium is ${}_kV({}_n\ddot{a}_x)$, that equals ${}_n\ddot{a}_x - \bar{P}({}_n\ddot{a}_x)\ddot{a}_{x+k:\overline{n-k}|}$ when $x \dots n$, \ddot{a}_{x+k} otherwise. The figure is shown in 1.




Figure 1: Benefit reserve of \ddot{a}_{65}

Insurance and annuities on two heads

Lifecontingencies package provides functions to evaluate life insurance and annuities on two lives. Following examples will check the equality $a_{\overline{xy}} = a_x + a_y - a_{xy}$.

```
R> axn(soa08Act, x=65,m=1)+axn(soa08Act, x=70,m=1)-axyn(soa08Act,soa08Act, x=65,y=
```

```
[1] 10.35704
```

```
R> axyn(soa08Act,soa08Act, x=65,y=70, status="last",m=1)
```

```
[1] 10.35704
```

Reversionary annuity (annuities payable to life y upon death of x), $a_{x|y} = a_y - a_{xy}$ can also be evaluate using **lifecontingencies** functions.

```
R> #assume x aged 65, y aged 60
```

```
R> axn(soa08Act, x=60,m=1)-axyn(soa08Act,soa08Act, x=65,y=60,status="joint",m=1)
```

```
[1] 2.695232
```

4.4. Stochastic analysis

This last paragraphs will show some stochastic analysis that can be performed by our package, both in demographic analysis and life insurance evaluation.

The age-until-death, both in the continuous (T_x) or curtate form (K_x), is a stochastic variable whose distribution is implicit within the deaths distribution of a given life table. The code below shows how to sample values from the age-until-death distribution implicit in the SOA life table.

```
R> data(soa08Act)
R> #sample 10 numbers from the Tx distribution
R> sample1<-rLife(n=10,object=soa08Act,x=0,type="Tx")
R> #sample 10 numbers from the Kx distribution
R> sample1<-rLife(n=10,object=soa08Act,x=0,type="Kx")
```

while code below shows how the mean of the sampled distribution is statistically equivalent to the expected life time.

```
R>          #assume an insured aged 29
R>          #his expected integer number of years until death is
R>          exn(soa08Act, x=29,type="curtate")

[1] 45.50066

R>          #check if we are sampling from a statistically equivalent distribution
R>          t.test(x=rLife(2000,soa08Act, x=29,type="Kx"),mu=exn(soa08Act, x=29,type="curtate"))

[1] 0.4193568

R>          #statistically not significant
```

Finally figure 2 shows the deaths distribution implicit in the ips55M life table.

The APV is a present value of a random variable that represents a composite function between the discount amount and indicator variables regarding the life status of the insured. Figure 3 shows the stochastic distribution of \ddot{a}_{65} .

5. Discussion

The **lifecontingencies** package allows actuaries to perform financial mathematics and life contingencies actuarial mathematics within R. It offers the basic tools to manipulate life tables and perform financial calculations. Pricing, reserving and stochastic evaluations of most important life insurance contract can be performed within R.

One of the most important limitations of **lifecontingencies** is handling only single decrement tables. In the future the **lifetable** class will probably be expanded to handle multiple decrement causes. Moreover in the future we expect to provide coerce methods toward packages




Figure 2: Deaths distribution implicit in the IPS55 males table

specialized in demographic analysis, like **demography** and **LifeTables**. Communciation with interest rates modelling packages, as **termstrcR** will be also explored.

Disclaimer

The accuracy of calculation have been verified by checkings with numerical examples reported in [Bowers *et al.* \(1997\)](#). The package numerical results are identical to those reported in the [Bowers *et al.* \(1997\)](#) for most function, with the exception of fractional payments annuities where the accuracy leads only to the 5th decimal. The reason of such inaccuracy is due to the fact that the package calculates the APV by directly sum of fractional survival probabilities, while the formulas reported in [Bowers *et al.* \(1997\)](#) uses an analytical formula.

Acknowledgments

I wish to thank Christophe Dutang and Tim Riffe for their valuable suggestions.




Figure 3: Stochastic distribution of \ddot{a}_{65}

References

- Anderson J, of Actuaries Education S, Committee E (1999). *Commutation functions*. Education and Examination Committee of the Society of Actuaries.
- Bowers N, Gerber H, Hickman J, Jones D, Nesbitt C (1997). “Actuarial Mathematics. Schaumburg.” *IL: Society of Actuaries*, pp. 79–82.
- Broverman S (2008). *Mathematics of investment and credit*. ACTEX academic series. ACTEX Publications. ISBN 9781566986571.
- Chambers J (2008). *Software for data analysis: programming with R*. Statistics and computing. Springer. ISBN 9780387759357. URL <http://books.google.com/books?id=UXneu0IvhEAC>.
- Dutang C, Goulet V, Pigeon M (2008). “actuar: An R Package for Actuarial Science.” *Journal of Statistical Software*, **25**(7), 38. URL <http://www.jstatsoft.org/v25/i07>.
- Ferstl R, Hayden J (2010). “Zero-Coupon Yield Curve Estimation with the Package termstrc.” *Journal of Statistical Software*, **36**(1), 1–34. URL <http://www.jstatsoft.org/v36/i01/>.
- Gesmann M, Zhang Y (2011). *ChainLadder: Mack, Bootstrap, Munich and Multivariate-chain-ladder Methods*. R package version 0.1.4-3.4.
- Guirrieri S (2010). *Simulating the Term Structure of Interest Rates with arbitrary marginals*. Ph.D. thesis, University of Palermo - Department of Statistics and Mathematics ”S. Vianelli”, Palermo. URL <http://www.guirrieri.host22.com>.
- Keyfitz N, Caswell H (2005). *Applied mathematical demography*. Statistics for biology and health. Springer. ISBN 9780387225371. URL <http://books.google.it/books?id=PxsVxES7Sj0C>.
- Klugman S, Panjer H, Willmot G, Venter G (2009). *Loss models: from data to decisions*. Third edition. Wiley New York.
- Riffe T (2011). *LifeTable: LifeTable, a package with a small set of useful lifetable functions*. R package version 1.0.1, URL <http://sites.google.com/site/timriffepersonal/r-code/lifeable>.
- Rigby RA, Stasinopoulos DM (2005). “Generalized additive models for location, scale and shape,(with discussion).” *Applied Statistics*, **54**, 507–554.
- Rob J Hyndman, Heather Booth, Leonie Tickle, John Maindonald (2011). *demography: Forecasting mortality, fertility, migration and population data*. R package version 1.09-1, URL <http://CRAN.R-project.org/package=demography>.
- Ruckman C, Francis J (????). “FINANCIAL MATHEMATICS:A Practical Guide for Actuaries and other Business Professionals.”

Affiliation:

Giorgio Alfredo Spedicato

StatisticalAdvisor

Via Firenze 11 20037 Italy

Telephone: +39/334/6634384

E-mail: lifecontingencies@statisticaladvisor.com

URL: www.statisticaladvisor.com

Draft